



Benefit Investigation Request and Prescription Form

Phone: 855-802-8746
 Fax: 855-454-8746
 MyQUTENZAConnect.com
 Hours: (M-F) 9 AM-7 PM ET

Case ID: _____
 Received: _____

PATIENT INFORMATION					
Last Name	First Name	MI	Sex Assignment ¹ <input type="checkbox"/> Male <input type="checkbox"/> Female	Date of Birth	Phone
Address	City	State	ZIP	Email	
Patient's Initial Treatment?	<input type="checkbox"/> Yes <input type="checkbox"/> No - If No, Date of Initial Treatment:		Allergies	Anticipated Treatment Date	

MEDICAL INSURANCE - PRIMARY	
Plan Name	Phone
Member ID	Group #

MEDICAL INSURANCE - SECONDARY	
Plan Name	Phone
Member ID	Group #

PHARMACY INSURANCE - PRIMARY	
Member ID	BIN
PCN	Group #

PHARMACY INSURANCE - SECONDARY	
Member ID	BIN
PCN	Group #

PRESCRIBER INFORMATION			
Prescriber's Full Name	Practice Name	Practice Contact	
Address	City	State	ZIP
Phone	Fax	NPI Number	Tax ID Number

CLINICAL INFORMATION		
ICD-10-CM Code	CPT Code	A list of codes that may be appropriate can be found in the Access Tool Kit. It is the physician's responsibility to provide the correct indication and codes.
<input type="checkbox"/> Postherpetic Neuralgia (PHN)	<input type="checkbox"/> Diabetic Peripheral Neuropathy (DPN) of the Feet	<input type="checkbox"/> Other:
<input type="checkbox"/> Physician Office	<input type="checkbox"/> Hospital Outpatient	<input type="checkbox"/> Other Site of Service:

PRESCRIPTION INFORMATION			
	Quantity # of Topical Systems (280 cm ² billing units) _____	Refills _____	Fulfillment Options <input type="checkbox"/> 1 Kit (carton includes 1 topical system and Cleansing Gel) NDC #72512-928-01 <input type="checkbox"/> 2 Kit (carton includes 2 topical systems and Cleansing Gel) NDC #72512-929-01 <input type="checkbox"/> 4 Kit (carton includes 4 topical systems and Cleansing Gel) NDC #72512-930-01
	HAIC Levels <input type="checkbox"/> Auto Transfer By checking this box, if Rx coverage is found, your prescription will be automatically transferred to a specialty pharmacy for fulfillment.	Directions	Shipping Address (if different from above)

ATTACH THE PATIENT CHART AND/OR CLINICAL DATA WITH THE SUBMISSION OF THIS INTAKE FORM TO BEGIN THE PA PROCESS.

PRESCRIBER'S SIGNATURE ²	
<input type="checkbox"/> Automatically re-investigate patient for potential retreatment in 91 days	
Prescriber's Signature: _____	Date: _____

- Gender override edits may be permissible by payer.
- Authorization for Release of Health Information: By signing this form, I represent to My QUTENZA Connect that I have obtained all necessary federal and state authorizations and consents from my patient to allow me to release health information to My QUTENZA Connect and its contracted third parties. I authorize My QUTENZA Connect to act on my behalf for the limited purposes of transmitting this prescription to the appropriate pharmacy designated by the patient utilizing their benefit plan. My signature on this form also provides consent to contact this patient's insurance provider for this prescription on the prescriber's behalf.





Benefit Investigation Request and Prescription Form

Phone: 855-802-8746
Fax: 855-454-8746
MyQUTENZAConnect.com
Hours: (M-F) 9 AM–7 PM ET

INDICATION

QUTENZA® (capsaicin) 8% topical system is indicated in adults for the treatment of neuropathic pain associated with postherpetic neuralgia (PHN) or associated with diabetic peripheral neuropathy (DPN) of the feet.

IMPORTANT SAFETY INFORMATION

Do not dispense QUTENZA to patients for self-administration or handling. Use only on dry, unbroken skin. Only physicians or healthcare professionals are to administer and handle QUTENZA, following the procedures in the label.

WARNINGS AND PRECAUTIONS

- **Severe Irritation:** Whether applied directly or transferred accidentally from other surfaces, capsaicin can cause severe irritation of eyes, mucous membranes, respiratory tract, and skin to the healthcare professional, patients, and others. Do not use near eyes or mucous membranes, including face and scalp. Take protective measures, including wearing nitrile gloves and not touching items or surfaces that the patient may also touch. Flush irritated mucous membranes or eyes with water and provide supportive medical care for shortness of breath. Remove affected individuals from the vicinity of QUTENZA. Do not re-expose affected individuals to QUTENZA if respiratory irritation worsens or does not resolve. If skin not intended to be treated comes into contact with QUTENZA, apply Cleansing Gel and then wipe off with dry gauze. Thoroughly clean all areas and items exposed to QUTENZA and dispose of properly. Because aerosolization of capsaicin can occur with rapid removal, administer QUTENZA in a well-ventilated area, and remove gently and slowly, rolling the adhesive side inward.
- **Application-Associated Pain:** Patients may experience substantial procedural pain and burning upon application and following removal of QUTENZA. Prepare to treat acute pain during and following application with local cooling and/or appropriate analgesic medication.

- **Increase in Blood Pressure:** Transient increases in blood pressure may occur with QUTENZA treatment. Monitor blood pressure during and following treatment procedure and provide support for treatment-related pain. Patients with unstable or poorly controlled hypertension, or a recent history of cardiovascular or cerebrovascular events, may be at an increased risk of adverse cardiovascular effects. Consider these factors prior to initiating QUTENZA treatment.
- **Sensory Function:** Reductions in sensory function (generally minor and temporary) have been reported following administration of QUTENZA. Assess for signs of sensory deterioration or loss prior to each application of QUTENZA. If sensory loss occurs, treatment should be reconsidered.
- **Severe Application Site Burns:** Full-thickness (third-degree) and deep partial-thickness (second-degree) burns have been reported following administration of QUTENZA. Cases of full-thickness (third-degree) burns, requiring hospitalization and skin grafting have been reported in patients who received QUTENZA for an unapproved indication and/or frequency of dosing at an application site where there had been prior skin trauma. Ensure that dosage and administration recommendations are followed.

ADVERSE REACTIONS

The most common adverse reactions ($\geq 5\%$ and $>$ control group) in all controlled clinical trials are application site erythema, application site pain, and application site pruritus.

To report SUSPECTED ADVERSE REACTIONS, contact Averitas Pharma, Inc. at 1-877-900-6479 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see accompanying [full Prescribing Information](#).